

# RT<sup>2</sup> Profiler PCR Array (96-Well Format and 384-Well [4 x 96] Format)

## Human GABA & Glutamate

Cat. no. 330231 PAHS-152ZA

For pathway expression analysis

Format	For use with the following real-time cyclers
RT <sup>2</sup> Profiler PCR Array, Format A	Applied Biosystems <sup>®</sup> models 5700, 7000, 7300, 7500, 7700, 7900HT, ViiA™ 7 (96-well block); Bio-Rad <sup>®</sup> models iCycler <sup>®</sup> , iQ™ 5, MyiQ™, MyiQ2; Bio-Rad/MJ Research Chromo4™; Eppendorf <sup>®</sup> Mastercycler <sup>®</sup> ep realplex models 2, 2s, 4, 4s; Stratagene <sup>®</sup> models Mx3005P <sup>®</sup> , Mx3000P <sup>®</sup> ; Takara TP-800
RT <sup>2</sup> Profiler PCR Array, Format C	Applied Biosystems models 7500 (Fast block), 7900HT (Fast block), StepOnePlus™, ViiA 7 (Fast block)
RT <sup>2</sup> Profiler PCR Array, Format D	Bio-Rad CFX96™; Bio-Rad/MJ Research models DNA Engine Opticon <sup>®</sup> , DNA Engine Opticon 2; Stratagene Mx4000 <sup>®</sup>
RT <sup>2</sup> Profiler PCR Array, Format E	Applied Biosystems models 7900HT (384-well block), ViiA 7 (384-well block); Bio-Rad CFX384™
RT <sup>2</sup> Profiler PCR Array, Format F	Roche <sup>®</sup> LightCycler <sup>®</sup> 480 (96-well block)
RT <sup>2</sup> Profiler PCR Array, Format G	Roche LightCycler 480 (384-well block)
RT <sup>2</sup> Profiler PCR Array, Format H	Fluidigm <sup>®</sup> BioMark™



Sample & Assay Technologies

## Description

The Human GABA & Glutamate RT<sup>2</sup> Profiler PCR Array profiles the expression of 84 key genes of the GABA (γ-aminobutyric acid) and glutamate neurotransmitter systems. The brain is a complex organ, capable of exerting essential responses for a variety of internal and external stimuli. Each stimulus activates excitatory or inhibitory responses, and the sum of these responses on each neuron results in propagated or inhibited neuronal transmission. Of the wide variety of neuronal receptors in the brain, the major excitatory receptors recognize the ligand glutamate, and the major inhibitory receptors respond to the ligand GABA. The GABA neurotransmitter system includes the GABAA and GABAC classes of ligand gated ion channels. The glutamate neurotransmitter system includes NMDA, AMPA, and kainate ligand-gated ion channels. Key enzymes synthesize GABA or glutamate as necessary, which are then transported into synaptic vesicles. Release of GABA or glutamate from vesicles activates postsynaptic GABA-responsive or glutamate-responsive ion channels, respectively, initiating downstream G protein signaling to propagate neurotransmission. Dysregulation of GABAergic or glutamatergic synaptic transmission results in a wide variety of nervous system disorders, including chronic pain, psychiatric diseases, neurodegenerative diseases, and insomnia. There are many drugs that are agonists or antagonists of the GABA and glutamate neurotransmitter systems. Historically, these major excitatory and inhibitory systems were studied separately. However, recent research suggests that input from both GABA and glutamate are necessary for normal nervous system growth and function. This array represents genes essential for the synthesis and transport of GABA and glutamate, as well as responsive ion channels and downstream signaling. The results of this array may yield insights into the interaction of these excitatory and inhibitory neuronal systems during essential cognitive functions. Using real-time PCR, research studies can easily and reliably analyze the expression of a focused panel of genes involved in neuronal GABA and glutamate functions with this array.

For further details, consult the *RT<sup>2</sup> Profiler PCR Array Handbook*.

## Shipping and storage

RT<sup>2</sup> Profiler PCR Arrays in formats A, C, D, E, F, and G are shipped at ambient temperature, on dry ice, or blue ice packs depending on destination and accompanying products. RT<sup>2</sup> Profiler PCR Arrays in format H are shipped on dry ice or blue ice packs.

For long term storage, keep plates at –20°C.

**Note:** Ensure that you have the correct RT<sup>2</sup> Profiler PCR Array format for your real-time cycler (see table above).

**Note:** Open the package and store the products appropriately immediately on receipt.

---

## Array layout (96-well)

For 384-well 4 x 96 PCR arrays, genes are present in a staggered format. Refer to the *RT<sup>2</sup> Profiler PCR Array Handbook* for layout.

	1	2	3	4	5	6	7	8	9	10	11	12
A	ABAT	ADCY7	ADORA1	ADORA2A	ALDH5A1	APP	AVP	BDNF	CACNA1A	CACNA1B	CDK5R1	CLN3
B	DLG4	GABBR1	GABBR2	GABRA1	GABRA2	GABRA4	GABRA5	GABRA6	GABRB1	GABRB3	GABRD	GABRE
C	GABRG1	GABRG2	GABRG3	GABRQ	GABRR1	GABRR2	GAD1	GLS	GLUL	GNAI1	GNAQ	GPHN
D	GRIA1	GRIA2	GRIA3	GRIA4	GRIK1	GRIK2	GRIK4	GRIK5	GRIN1	GRIN2A	GRIN2B	GRIN2C
E	GRM1	GRM2	GRM3	GRM4	GRM5	GRM6	GRM7	GRM8	HOMER1	HOMER2	IL1B	ITPR1
F	MAPK1	NSF	P2RX7	PHGDH	PLA2G6	PLCB1	PRODH	SHANK2	SLC17A6	SLC17A7	SLC17A8	SLC1A1
G	SLC1A2	SLC1A3	SLC1A6	SLC32A1	SLC38A1	SLC6A1	SLC6A11	SLC6A12	SLC6A13	SLC7A11	SNCA	SRR
H	ACTB	B2M	GAPDH	HPRT1	RPLP0	HGDC	RTC	RTC	RTC	PPC	PPC	PPC

## Gene table: RT<sup>2</sup> Profiler PCR Array

Position	UniGene	GenBank	Symbol	Description
A01	Hs.336768	NM_000663	ABAT	4-aminobutyrate aminotransferase
A02	Hs.513578	NM_001114	ADCY7	Adenylate cyclase 7
A03	Hs.77867	NM_000674	ADORA1	Adenosine A1 receptor
A04	Hs.197029	NM_000675	ADORA2A	Adenosine A2a receptor
A05	Hs.371723	NM_001080	ALDH5A1	Aldehyde dehydrogenase 5 family, member A1
A06	Hs.434980	NM_000484	APP	Amyloid beta (A4) precursor protein
A07	Hs.89648	NM_000490	AVP	Arginine vasopressin
A08	Hs.502182	NM_001709	BDNF	Brain-derived neurotrophic factor
A09	Hs.501632	NM_000068	CACNA1A	Calcium channel, voltage-dependent, P/Q type, alpha 1A subunit
A10	Hs.495522	NM_000718	CACNA1B	Calcium channel, voltage-dependent, N type, alpha 1B subunit
A11	Hs.500015	NM_003885	CDK5R1	Cyclin-dependent kinase 5, regulatory subunit 1 (p35)
A12	Hs.628393	NM_000086	CLN3	Ceroid-lipofuscinosis, neuronal 3
B01	Hs.463928	NM_001365	DLG4	Discs, large homolog 4 (Drosophila)
B02	Hs.167017	NM_001470	GABBR1	Gamma-aminobutyric acid (GABA) B receptor, 1
B03	Hs.198612	NM_005458	GABBR2	Gamma-aminobutyric acid (GABA) B receptor, 2
B04	Hs.175934	NM_000806	GABRA1	Gamma-aminobutyric acid (GABA) A receptor, alpha 1
B05	Hs.116250	NM_000807	GABRA2	Gamma-aminobutyric acid (GABA) A receptor, alpha 2
B06	Hs.248112	NM_000809	GABRA4	Gamma-aminobutyric acid (GABA) A receptor, alpha 4
B07	Hs.612087	NM_000810	GABRA5	Gamma-aminobutyric acid (GABA) A receptor, alpha 5
B08	Hs.90791	NM_000811	GABRA6	Gamma-aminobutyric acid (GABA) A receptor, alpha 6
B09	Hs.27283	NM_000812	GABRB1	Gamma-aminobutyric acid (GABA) A receptor, beta 1
B10	Hs.302352	NM_000814	GABRB3	Gamma-aminobutyric acid (GABA) A receptor, beta 3
B11	Hs.113882	NM_000815	GABRD	Gamma-aminobutyric acid (GABA) A receptor, delta
B12	Hs.22785	NM_004961	GABRE	Gamma-aminobutyric acid (GABA) A receptor, epsilon
C01	Hs.375051	NM_173536	GABRG1	Gamma-aminobutyric acid (GABA) A receptor, gamma 1
C02	Hs.7195	NM_000816	GABRG2	Gamma-aminobutyric acid (GABA) A receptor, gamma 2
C03	Hs.569475	NM_033223	GABRG3	Gamma-aminobutyric acid (GABA) A receptor, gamma 3
C04	Hs.283081	NM_018558	GABRRQ	Gamma-aminobutyric acid (GABA) receptor, theta
C05	Hs.437745	NM_002042	GABRR1	Gamma-aminobutyric acid (GABA) receptor, rho 1
C06	Hs.99927	NM_002043	GABRR2	Gamma-aminobutyric acid (GABA) receptor, rho 2
C07	Hs.420036	NM_000817	GAD1	Glutamate decarboxylase 1 (brain, 67kDa)
C08	Hs.116448	NM_014905	GLS	Glutaminase
C09	Hs.518525	NM_002065	GLUL	Glutamate-ammonia ligase
C10	Hs.134587	NM_002069	GNAI1	Guanine nucleotide binding protein (G protein), alpha inhibiting activity polypeptide 1
C11	Hs.269782	NM_002072	GNAQ	Guanine nucleotide binding protein (G protein), q polypeptide
C12	Hs.208765	NM_020806	GPHN	Gephyrin
D01	Hs.519693	NM_000827	GRIA1	Glutamate receptor, ionotropic, AMPA 1
D02	Hs.32763	NM_000826	GRIA2	Glutamate receptor, ionotropic, AMPA 2
D03	Hs.377070	NM_000828	GRIA3	Glutamate receptor, ionotropic, AMPA 3
D04	Hs.503743	NM_000829	GRIA4	Glutamate receptor, ionotropic, AMPA 4
D05	Hs.706747	NM_000830	GRIK1	Glutamate receptor, ionotropic, kainate 1
D06	Hs.98262	NM_021956	GRIK2	Glutamate receptor, ionotropic, kainate 2
D07	Hs.568901	NM_014619	GRIK4	Glutamate receptor, ionotropic, kainate 4
D08	Hs.367799	NM_002088	GRIK5	Glutamate receptor, ionotropic, kainate 5

Position	UniGene	GenBank	Symbol	Description
D09	Hs.558334	NM_007327	GRIN1	Glutamate receptor, ionotropic, N-methyl D-aspartate 1
D10	Hs.411472	NM_000833	GRIN2A	Glutamate receptor, ionotropic, N-methyl D-aspartate 2A
D11	Hs.654430	NM_000834	GRIN2B	Glutamate receptor, ionotropic, N-methyl D-aspartate 2B
D12	Hs.436980	NM_000835	GRIN2C	Glutamate receptor, ionotropic, N-methyl D-aspartate 2C
E01	Hs.32945	NM_000838	GRM1	Glutamate receptor, metabotropic 1
E02	Hs.121510	NM_000839	GRM2	Glutamate receptor, metabotropic 2
E03	Hs.590575	NM_000840	GRM3	Glutamate receptor, metabotropic 3
E04	Hs.654847	NM_000841	GRM4	Glutamate receptor, metabotropic 4
E05	Hs.147361	NM_000842	GRM5	Glutamate receptor, metabotropic 5
E06	Hs.248131	NM_000843	GRM6	Glutamate receptor, metabotropic 6
E07	Hs.606393	NM_000844	GRM7	Glutamate receptor, metabotropic 7
E08	Hs.449625	NM_000845	GRM8	Glutamate receptor, metabotropic 8
E09	Hs.591761	NM_004272	HOMER1	Homer homolog 1 (Drosophila)
E10	Hs.578443	NM_004839	HOMER2	Homer homolog 2 (Drosophila)
E11	Hs.126256	NM_000576	IL1B	Interleukin 1, beta
E12	Hs.567295	NM_002222	ITPR1	Inositol 1,4,5-trisphosphate receptor, type 1
F01	Hs.431850	NM_002745	MAPK1	Mitogen-activated protein kinase 1
F02	Hs.431279	NM_006178	NSF	N-ethylmaleimide-sensitive factor
F03	Hs.729169	NM_002562	P2RX7	Purinergic receptor P2X, ligand-gated ion channel, 7
F04	Hs.487296	NM_006623	PHGDH	Phosphoglycerate dehydrogenase
F05	Hs.170479	NM_003560	PLA2G6	Phospholipase A2, group VI (cytosolic, calcium-independent)
F06	Hs.431173	NM_015192	PLCB1	Phospholipase C, beta 1 (phosphoinositide-specific)
F07	Hs.517352	NM_016335	PRODH	Proline dehydrogenase (oxidase) 1
F08	Hs.268726	NM_133266	SHANK2	SH3 and multiple ankyrin repeat domains 2
F09	Hs.242821	NM_020346	SLC17A6	Solute carrier family 17 (sodium-dependent inorganic phosphate cotransporter), member 6
F10	Hs.375616	NM_020309	SLC17A7	Solute carrier family 17 (sodium-dependent inorganic phosphate cotransporter), member 7
F11	Hs.116871	NM_139319	SLC17A8	Solute carrier family 17 (sodium-dependent inorganic phosphate cotransporter), member 8
F12	Hs.444915	NM_004170	SLC1A1	Solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system Xag), member 1
G01	Hs.502338	NM_004171	SLC1A2	Solute carrier family 1 (glial high affinity glutamate transporter), member 2
G02	Hs.481918	NM_004172	SLC1A3	Solute carrier family 1 (glial high affinity glutamate transporter), member 3
G03	Hs.515217	NM_005071	SLC1A6	Solute carrier family 1 (high affinity aspartate/glutamate transporter), member 6
G04	Hs.179080	NM_080552	SLC32A1	Solute carrier family 32 (GABA vesicular transporter), member 1
G05	Hs.533770	NM_030674	SLC38A1	Solute carrier family 38, member 1
G06	Hs.443874	NM_003042	SLC6A1	Solute carrier family 6 (neurotransmitter transporter, GABA), member 1
G07	Hs.657405	NM_014229	SLC6A11	Solute carrier family 6 (neurotransmitter transporter, GABA), member 11
G08	Hs.437174	NM_003044	SLC6A12	Solute carrier family 6 (neurotransmitter transporter, betaine/GABA), member 12
G09	Hs.504398	NM_016615	SLC6A13	Solute carrier family 6 (neurotransmitter transporter, GABA), member 13
G10	Hs.390594	NM_014331	SLC7A11	Solute carrier family 7 (anionic amino acid transporter light chain, xc- system), member 11
G11	Hs.271771	NM_000345	SNCA	Synuclein, alpha (non A4 component of amyloid precursor)
G12	Hs.461954	NM_021947	SRR	Serine racemase
H01	Hs.520640	NM_001101	ACTB	Actin, beta
H02	Hs.534255	NM_004048	B2M	Beta-2-microglobulin
H03	Hs.592355	NM_002046	GAPDH	Glyceraldehyde-3-phosphate dehydrogenase
H04	Hs.412707	NM_000194	HPRT1	Hypoxanthine phosphoribosyltransferase 1
H05	Hs.546285	NM_001002	RPLP0	Ribosomal protein, large, P0
H06	N/A	SA_00105	HGDC	Human Genomic DNA Contamination
H07	N/A	SA_00104	RTC	Reverse Transcription Control
H08	N/A	SA_00104	RTC	Reverse Transcription Control
H09	N/A	SA_00104	RTC	Reverse Transcription Control
H10	N/A	SA_00103	PPC	Positive PCR Control
H11	N/A	SA_00103	PPC	Positive PCR Control
H12	N/A	SA_00103	PPC	Positive PCR Control

## Related products

For optimal performance, RT<sup>2</sup> Profiler PCR Arrays should be used together with the RT<sup>2</sup> First Strand Kit for cDNA synthesis and RT<sup>2</sup> SYBR<sup>®</sup> Green qPCR Mastermixes for PCR.

Product	Contents	Cat. no.
RT <sup>2</sup> First Strand Kit (12)	Enzymes and reagents for cDNA synthesis	330401
RT <sup>2</sup> SYBR Green qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with real-time cyclers that do not require a reference dye, including: Bio-Rad models CFX96, CFX384, DNA Engine Opticon 2; Bio-Rad/MJ Research Chromo4; Roche LightCycler 480 (96-well and 384-well); all other cyclers	330500
RT <sup>2</sup> SYBR Green ROX™ qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Applied Biosystems models 5700, 7000, 7300, 7500 [Standard and FAST], 7700, 7900HT 96-well block [Standard and FAST] and 384-well block, StepOnePlus; Eppendorf Mastercycler ep realplex models 2, 2S, 4, 4S; Stratagene models Mx3000P, Mx3005P, Mx4000; Takara TP-800	330520
RT <sup>2</sup> SYBR Green Fluor qPCR Mastermix (2)*	For 2 x 96 assays in 96-well plates; suitable for use with the following real-time cyclers: Bio-Rad models iCycler, iQ5, MyiQ, MyiQ2	330510

\* Larger kit sizes available; please inquire.

RT<sup>2</sup> Profiler PCR Array products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

For up-to-date licensing information and product-specific disclaimers, see the respective QIAGEN kit handbook or user manual. QIAGEN kit handbooks and user manuals are available at [www.qiagen.com](http://www.qiagen.com) or can be requested from QIAGEN Technical Services or your local distributor.

Trademarks: QIAGEN® (QIAGEN Group); Applied Biosystems®, ViiA™, StepOnePlus™, ROX™ (Applied Biosystems Corporation or its subsidiaries); Bio-Rad®, iCycler®, iQ™, MyiQ™, Chromo4™, CFX96™, DNA Engine Opticon®, CFX384™ (Bio-Rad Laboratories, Inc.); Stratagene®, Mx3005P®, Mx3000P®, Mx4000® (Stratagene); Eppendorf®, Mastercycler® (Eppendorf AG); Roche®, LightCycler® (Roche Group); Fluidigm® BioMark™ (Fluidigm Corporation); SYBR® (Molecular Probes, Inc.).

1066029 03/2011 © 2011 QIAGEN, all rights reserved.

[www.qiagen.com](http://www.qiagen.com)

Canada ■ 800-572-9613

Ireland ■ 1800 555 049

Norway ■ 800-18859

China ■ 8621-3865-3865

Italy ■ 800-787980

Singapore ■ 1800-742-4368

Denmark ■ 80-885945

Japan ■ 03-6890-7300

Spain ■ 91-630-7050

Australia ■ 1-800-243-800

Finland ■ 0800-914416

Korea (South) ■ 080-000-7145

Sweden ■ 020-790282

Austria ■ 0800/281010

France ■ 01-60-920-930

Luxembourg ■ 8002 2076

Switzerland ■ 055-254-22-11

Belgium ■ 0800-79612

Germany ■ 02103-29-12000

Mexico ■ 01-800-7742-436

UK ■ 01293-422-911

Brazil ■ 0800-557779

Hong Kong ■ 800 933 965

The Netherlands ■ 0800 0229592

USA ■ 800-426-8157



Sample & Assay Technologies